

Exam 2

1. Question

Given the following information:

$$\text{🍍} + \text{🍍} + \text{🍌} = 674$$

$$\text{🍌} + \text{🍌} + \text{🍍} = 385$$

$$\text{🍌} + \text{🍊} + \text{🍌} = 102$$

Compute:

$$\text{🍌} + \text{🍊} + \text{🍍} = ?$$

- a. 321
- b. 391
- c. 568
- d. 561
- e. 385

Solution

The information provided can be interpreted as the price for three fruit baskets with different combinations of the three fruits. This corresponds to a system of linear equations where the price of the three fruits is the vector of unknowns x :

$$x_1 = \text{🍌} \quad x_2 = \text{🍊} \quad x_3 = \text{🍍}$$

The system of linear equations is then:

$$\begin{pmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \\ 2 & 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 674 \\ 385 \\ 102 \end{pmatrix}$$

This can be solved using any solution algorithm, e.g., elimination:

$$x_1 = 32, x_2 = 38, x_3 = 321.$$

Based on the three prices for the different fruits it is straightforward to compute the total price of the fourth fruit basket via:

$$\begin{array}{ccccccc} \text{🍌} & + & \text{🍊} & + & \text{🍍} & = & \\ x_1 & + & x_2 & + & x_3 & = & \end{array}$$

$$32 + 38 + 321 = 391$$

- a. False
- b. True
- c. False
- d. False
- e. False

2. Question


What is the distance between the two points $p = (1, 2)$ and $q = (5, 4)$ in a Cartesian coordinate system?

- a. 2.971
- b. 4.472
- c. 1.118
- d. 3.888
- e. 2.449

Solution

The distance d of p and q is given by $d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(1 - 5)^2 + (2 - 4)^2} = 4.472$.



- a. False
- b. True
- c. False
- d. False
- e. False

3. Question

What is the derivative of $f(x) = x^8 e^{2.7x}$, evaluated at $x = 0.53$?

Solution

Using the product rule for $f(x) = g(x) \cdot h(x)$, where $g(x) := x^8$ and $h(x) := e^{2.7x}$, we obtain

$$\begin{aligned} f'(x) &= [g(x) \cdot h(x)]' = g'(x) \cdot h(x) + g(x) \cdot h'(x) \\ &= 8x^{8-1} \cdot e^{2.7x} + x^8 \cdot e^{2.7x} \cdot 2.7 \\ &= e^{2.7x} \cdot (8x^7 + 2.7x^8) \\ &= e^{2.7x} \cdot x^7 \cdot (8 + 2.7x). \end{aligned}$$

Evaluated at $x = 0.53$, the answer is

$$e^{2.7 \cdot 0.53} \cdot 0.53^7 \cdot (8 + 2.7 \cdot 0.53) = 0.463409.$$

Thus, rounded to two digits we have $f'(0.53) = 0.46$.

4. Question

The daily expenses of summer tourists in Vienna are analyzed. A survey with 112 tourists is conducted. This shows that the tourists spend on average 126 EUR. The sample variance s_{n-1}^2 is equal to 117.

Determine a 95% confidence interval for the average daily expenses (in EUR) of a tourist.

- What is the lower confidence bound?
- What is the upper confidence bound?

Solution

The 95% confidence interval for the average expenses μ is given by:

$$\begin{aligned} &\left[\bar{y} - 1.96 \sqrt{\frac{s_{n-1}^2}{n}}, \bar{y} + 1.96 \sqrt{\frac{s_{n-1}^2}{n}} \right] \\ &= \left[126 - 1.96 \sqrt{\frac{117}{112}}, 126 + 1.96 \sqrt{\frac{117}{112}} \right] \\ &= [123.997, 128.003]. \end{aligned}$$

- The lower confidence bound is 123.997.
- The upper confidence bound is 128.003.

5. Question

For 63 firms the number of employees X and the amount of expenses for continuing education Y (in EUR) were recorded. The statistical summary of the data set is given by:

	Variable X	Variable Y
Mean	51	202
Variance	95	1889

The correlation between X and Y is equal to 0.73.

Estimate the expected amount of money spent for continuing education by a firm with 52 employees using least squares regression.

Solution

First, the regression line $y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$ is determined. The regression coefficients are given by:

$$\hat{\beta}_1 = r \cdot \frac{s_y}{s_x} = 0.73 \cdot \sqrt{\frac{1889}{95}} = 3.2552,$$

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \cdot \bar{x} = 202 - 3.2552 \cdot 51 = 35.98504.$$

The estimated amount of money spent by a firm with 52 employees is then given by:

$$\hat{y} = 35.98504 + 3.2552 \cdot 52 = 205.255.$$